

## STEP 1: Mifepristone

Pill taken on \_\_\_\_\_ at \_\_\_\_\_ am/pm



Swallow 1 pill when you are ready.

## STEP 2: Misoprostol

Take pills on \_\_\_\_\_ at \_\_\_\_\_ am/pm



**Take these pills 24-48 hours after STEP 1**

Put 2 pills inside each cheek (4 pills total).

Wait 30 minutes for the pills to dissolve.

Then swish with water and swallow.



### **What should I expect?**

- Cramps and bleeding that may be worse than your normal period will start within a few hours of taking **STEP 2**.
- Bleeding might start after taking **STEP 1** and before taking **STEP 2**.
- You might have nausea, diarrhea, fever, or chills.
- Do not take any more pain pills than your doctor recommended. You can take an over-the-counter medicine for diarrhea, if needed.

### **When should I call my doctor or seek medical attention?**

- If you have bleeding that soaks 2 full-sized pads per hour for 2 hours in a row.
- If you have a fever over 101°F, or fever over 100.4 °F that lasts more than 4 hours.
- If you have pain that does not get better after taking pain medicine or severe pain in your belly or abdomen.
- General feelings of illness, weakness, or discomfort for more than 24 hours after **STEP 2**.
- If you have little bleeding or do not believe the pregnancy has passed.
- If you still have pregnancy symptoms 7 days after **STEP 2**.

**Your doctor may have you take misoprostol more than once. Follow your doctor's instructions.**